

TERNOVSKIY, P.F.

Effect of multiple scattering on the production of pairs of
charged particles in high energies in a medium. Zhur.eksp.i
teor.fiz. 37 no.4:1010-1016 O '59. (MIRA 13:5)

1. Moskovskiy gosudarstvennyy universitet.
(Electrons)

84720

17.1400
24.4500

S/056/60/039/001/040/041/XX
B006/B056

AUTHOR:

Ternovskiy, F. F.

TITLE:

Theory of Radiative Processes in Piecewise Homogeneous Media ²¹

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 39, No. 1(7), pp. 171-180

TEXT: L. D. Landau, I. Ya. Pomeranchuk, and A. B. Migdal (Refs. 1-4) have shown that radiative processes in media become much less intensive at high energies than what follows from the ordinary theory. M. L. Ter-Mikayelyan in this connection pointed out that, when investigating the radiation of the softest quanta, the effect of polarization of the medium must be taken into account. In the case of a finite medium, two types of transition radiation occur at the boundary. The first type is connected with a modification of the particle field due to polarization of the medium, and occurs at all energies and all rest masses of the particles. Here, an electromagnetic wave of less than optical frequency is emitted to the side that is opposite to the direction of motion. The spectrum of this radiation contains frequencies of up to $\omega = \omega_0 / \sqrt{1 - \beta^2}$, where

Card 1/3

84720

Theory of Radiative Processes in Piecewise
Homogeneous Media

S/056/60/039/001/040/041/XX
B006/B056

$\omega_0^2 = 4\pi nZe^2/m$, $v = \beta c$ is the particle velocity. The second type is connected with the influence of multiple scattering; it occurs only at sufficiently high energies, and can be observed only in the case of electrons and positrons. Its angular distribution has a marked maximum in the direction of motion, and the spectrum can, at sufficiently high energy, attain frequencies of the order of $mc^2/h\sqrt{1-\beta^2}$. A classical theory of this radiation was developed by I. I. Gol'dman (Ref. 9). In the present paper, the transition radiations of both types are quantum-theoretically investigated; it is first assumed that the semispace $z > 0$ is filled with a medium, for which $|1 - \sqrt{\epsilon}| \ll 1$, and $z < 0$ is assumed to be a vacuum. The radiation is investigated that occurs when a fast electron ($p_0 \gg 1$), coming from the vacuum, penetrates into the medium, ($\hbar = m = c = 1$), where multiple scattering and polarization of the medium are taken into account. In the following, the author investigates the radiation occurring when an electron passes through a plate of finite thickness. Using the results obtained by Migdal, the probabilities for bremsstrahlung and pair production by gamma quanta are calculated. The author thanks Professor A. B. Migdal for suggesting the problem and for discussions, V. D. Vaks for his help, and I. I. Gol'dman for making results available before their publication.

Card 2/3

84720

Theory of Radiative Processes in Piecewise
Homogeneous Media

S/056/60/039/001/040/041/XX
B006/B056

G. M. Caribyan is mentioned. There are 1 table and 11 references: 9 Soviet,
1 British, and 1 US.

ASSOCIATION: Moskovskiy gosudarstvennyy universitet
(Moscow State University)

SUBMITTED: February 25, 1960

Card 3/3

83201

S/056/60/039/002/038/044
B006/B070

24.6600
AUTHOR:

Ternovskiy, F. F.

TITLE:

The Effect of Multiple Scattering on the Proper Field of a Fast Charged Particle

PERIODICAL:

Zhurnal eksperimental'noy i teoreticheskoy fiziki, 1960,
Vol. 39, No. 2(8), pp. 491 - 496

TEXT: At first the author briefly discusses the related investigations of I. I. Gol'dman and G. M. Garibyan (Refs. 1,3). In the present paper, it is shown that the appearance of a transition radiation, that was predicted by the theory of Gol'dman, is related also to the change in the structure of the proper field of the particle, this change being a consequence of multiple scattering. This problem is treated in the second section of the paper, where the number of quanta per unit spectral interval are calculated. The number of quanta is equivalent to the proper field of the particles moving in the medium (Formula (6)). It is shown (Formula (9)) further, that the number of equivalent photons in vacuum, which are calculated according to the known formula of Weizsäcker and Williams

Card 1/3

83201

The Effect of Multiple Scattering on the
Proper Field of a Fast Charged Particle

S/056/60/039/002/038/044
B006/B070

(Ref. 4), agrees well with the number of transition quanta plus the number of photons which are equivalent to the field of the particle in the medium. For distances from the point of entrance in the medium that are smaller than the avalanche unit and in which the transition quanta cannot be absorbed, the cross section of direct pair formation is proportional to the number of equivalent photons in vacuum; for larger distances from the point of entrance which is no longer reached by transition radiation, the number of pairs directly produced by the field of the particle is proportional to the number of equivalent photons in the medium. Section 3 is concerned with this problem. In section 4, the author studies the process of pair formation by particles penetrating the medium for the case when the region, in which the radiation field is built, is of the same order of magnitude as the avalanche unit. Here it is not possible to consider separately the cross section for indirect pair formation. The total number of pairs formed in a depth which is large compared to the avalanche unit is given by the formula (19). A. B. Migdal, L. D. Landau, and I. Ya. Pomeranchuk are mentioned in the paper. There are 8 references: 7 Soviet and 1 US.

Card 2/3

The Effect of Multiple Scattering on the
Proper Field of a Fast Charged Particle

83201

S/056/60/039/002/038/044
B006/B070

ASSOCIATION: Moskovskiy gosudarstvennyy universitet (Moscow State
University)

SUBMITTED: March 25, 1960

4

Card 3/3

ACC NR: AP/002585

SOURCE CODE: UR/0020/66/171/005/1084/1087

AUTHOR: Ternovskiy, F. F.

ORG: Moscow State University im. M. V. Lomonosov (Moskovskiy gosudarstvennyy universitet); Scientific Research Institute of High Temperatures (Nauchno-issledovatel'skiy institut vysokikh temperatur)

TITLE: Certain effects due to diamagnetism of the winding of a superconducting solenoid

SOURCE: AN SSSR. Doklady, v. 171, no. 5, 1966, 1084-1087

TOPIC TAGS: superconductivity, solenoid, electric inductance, electric measurement, quantum effect

ABSTRACT: The purpose of the investigation was to ascertain what information concerning the properties of hard superconductors can be extracted from inductance measurements, since experiments have shown that the inductance of a superconductor depends greatly on the current flowing in it and on the external magnetic field. The properties of a superconducting solenoid are analyzed by using the dependence of the magnetic flux linking the turns of the solenoid on the current. The flux in turn is separated into external and internal components, with particular attention paid to the effect of the internal component, whose filamentary nature (quantum vortices) results in nonuniform magnetization, owing to the interaction between the quantum vortices and inhomogeneities in the superconducting material. An expression for the

Card 2/2

UDC: 537.312.62

ACC NR: AP7002385

internal flux is derived under certain assumptions concerning the distribution of the induction inside the solenoid, and a table of numerical values for the coefficients of this expression is presented. This leads to an expression for the inductance of the superconducting solenoid in the form of two components: external inductance, which is almost independent of the current, and internal inductance which depends both on the current and on the magnetic history of the solenoid. It is pointed out that usual impedance measurements give only the external inductance. It is noted that recent measurements of the internal flux in the superconductor agree with the approximate calculations presented in the paper. The author thanks V. V. Andrianov, V. B. Zenkevich, and V. V. Sychev for information on the results of the experiments and for fruitful discussions. This report was presented by Academician M. A. Leontovich 21 February 1966. Orig. art. has: 1 figure, 13 formulas, and 1 table.

SUB CODE: 2C/ SUBM DATE: 10Dec65/ ORIG REF: 002/ OTH REF: 006

Card 2/2

MELESHKIN, S.M., gornyy inzhener; BERLYAND, S.S., gornyy inzhener;
SIROTKIN, Z.L., inzh.; DENISOV, A.G., inzh.; TERNOVSKIY, G.I., inzh.;
BEKHTEREV, Yu.I., inzh.; ZOTOV, A.V., inzh.; IVANOV, E.I., inzh.;
VASIL'YEV, Ye.A., inzh.; SOLOV'YEVA, L.G., inzh.; D'YACHENKO, V.F.,
inzh.

Replies to V.V. Shan'ko's article "Efficient limits of using
truck haulage in open pits." Gor. zhur. no.1:75-77 Ja '62.

(MIRA 15:7)

1. Gosudarstvennyy nauchno-ekonomicheskii sovet Soveta Ministrov
SSSR (for Meleshkin). 2. Promtransproyekt Gosstroya SSSR (for
Berlyand). 3. Belorusskiy avtozavod (for Sirotkin, Denisov,
Ternovskiy, Bekhterev, Zotov, Ivanov). 4. Gosudarstvennyy
institut po proyektirovaniyu razrabotki rudnykh mestorozhdeniy
v yuzhnykh rayonov SSSR, Khar'kov (for Vasil'yev, Solov'yova,
D'yachenko).

(Mine haulage)
(Shan'ko, V.V.)

DENISOV, Aleksandr Gavrilovich; KAZAREZ, Aleksey Nikolayevich;
SIROTKIN, Zalya L'vovich; TERNOVSKIY, Genrikh Ivanovich;
SHUMSKIY, Mechislav Frantsevich; LESNYAKOV, F.I., red.;
GALAKTIONOVA, Ye.N., tekhn. red.

[MAZ-525 dump truck; its design and operation] Avtomobil'-
samosval MAZ-525; ustroistvo i ekspluatatsiia. Moskva,
Avtotransizdat, 1963. 166 p. (MIRA 16:10)
(Dump trucks)

TERNOVSKIY, G.V., Kapitan 1-go ranga v otstavke, Geroy Sovetskogo Soyuza

The feats of herats are immortal; on the 20th anniversary of
the liberation of Sevastopol. Mor. sbor. 47 no.5:19-22 My '64.
(MIRA 18:6)

1. 24233-65 EWT(m)
ACC NR: AI6014669

SOURCE CODE: UR/0241/65/010/010/0037/0041

AUTHOR: Yerokhin, R. A.--Erokhin, R. A.; Koshurnikova, N. A.; Ternovskiy, I. A.--
Ternovsky, I. A.

ORG: none

TITLE: Gamma-spectrometric intravital determination of Pu in the living organism

SOURCE: Meditsinskaya radiologiya, v. 10, no. 10, 1965, 37-41

TOPIC TAGS: plutonium, gamma spectrometer, radiology, americium, scintillation spectrometer, photomultiplier, pulse analyzer, pulse amplitude, rat, liver/FEU-24 photomultiplier, AI-100 pulse analyzer

ABSTRACT: The results of an experimental determination of the possibility of the direct intravital measurement of ^{239}Pu and Am^{241} in the organism by means of a scintillation gamma-spectrometer are presented. The principal components of the spectrometer used were: a NaI(Tl) scintillation crystal 20 mm thick and 40 mm in diameter, with an FEU-24 photomultiplier and an AI-100 pulse-amplitude analyzer. White rats were given, intravenously or intratracheally, Pu in the form of the nitrate salt $[\text{Pu}(\text{NO}_3)_4]$ with a pH value of 2.0, in the amount of 5 microcuries per rat, or Am^{241} in the form of the nitrate $[\text{Am}(\text{NO}_3)_3]$ with a pH of 2.8, in the amount of 2.72 microcuries per rat. The Pu and Am^{241} contents of the rat organism were measured immediately afterward as well as at intervals of 1, 2, 4, 8, 16, 32, and 64 days. Lung activity varied identically in rats intratracheally poisoned with Pu and Am^{241}

Cord 1/2

UDC: 616-008.927.994-073.584

L 24233-66

ACC NR: AP6014669

nitrites: toward the 16th day the activity dropped 30%; toward the 32nd day, 50%; and toward the end of the experiment, 65%. Throughout the experiment, the content of Pu exceeded that of Am²⁴¹. The dynamics of the change in liver activity following administration of the Pu nitrate toward the 3rd, 7th, 30th, and 60th day was 87, 72, 41.8, and 22.5%, respectively, for Am²⁴¹ these dynamics differed somewhat. In addition, the minimum amounts of Pu and Am²⁴¹ still detectable by the spectrometric method in different organs (kidneys, liver, hip, spleen) were determined; for Pu they were found to correspond to 0.019-0.045 microcuries per rat organ, and for Am²⁴¹, to 0.0028-0.0087 microcuries per rat organ. Orig. art. has: 2 figures and 4 tables. [JPRS]

SUB CODE: 06, 09, 20, 07 / SUBM DATE: 12Jan65 / ORIG REF: 003 / OTH REF: 005

Card 2/242

TERNOVSKIY, I. N.: "The use of sulfonin to treat cardiovascular."
 Min "ed in USSR. Kuzen' State Medical Academy in Nov.
 Krasnodar, 1966.
 (Dissertation for Degree of Candidate in Medical Sciences).

SO: Kulzhnaya literatura, No 31, 1966

TERNOVSKIY, I.N.

Dispensary services for mental patients in certain rural regions of Krasnodar Territory. Zhur. nevr. i psikh 58 no.12:1494-1497 '58.

(MIRA 12:1)

1. Kafedra psikhiatrii (zav. - prof. N.I. Bondarev) Kubanskogo meditsinskogo instituta.

(MENTAL DISORDERS, ther.
in rural cond. (Rus))

AUTHORS: Tulinova, V. B., Plyushchev, V. Ye., S/078/60/005/03/033/048
Ternovskaya, I. V., Lukova, S. N. B004/B005
Samuseva, R. G.

TITLE: Investigation of the Joint Solubility of Lanthanum and Sodium Sulfates


PERIODICAL: Zhurnal neorganicheskoy khimii, 1960, Vol 5, Nr 3, pp 695-700 (USSR)

ABSTRACT: The present paper is part of an extensive investigation of the physicochemical foundation of methods of separating rare earths which was started together with G. G. Urazov (Ref 3). The solubility in the system $\text{La}_2(\text{SO}_4)_3 - \text{Na}_2\text{SO}_4 - \text{H}_2\text{O}$ was determined at 25, 50, and 75°. The binary systems which are components of this system have been described in publications. The solubility was determined by the isothermal method. The equilibrium between solution and precipitate was established after 14 days which was checked analytically. The sulfate ion was determined gravimetrically as BaSO_4 , the lanthanum ion either gravimetrically as oxalate or, at low concentrations, colorimetrically according to reference 10. The results are shown in tables 1-3 (for 25, 50, and 75°), and as a diagram in figure 1. One double salt $\text{La}_2(\text{SO}_4)_3 \cdot \text{Na}_2\text{SO}_4 \cdot 2\text{H}_2\text{O}$ forms

Card 1/2

Investigation of the Joint Solubility of Lanthanum
and Sodium Sulfates

S/078/60/005/03/033/048
B004/B005

in the system investigated; its thermogram is shown in figure 2,
its Debye-pattern data in table 4. There are 1 figure, 4 tables,
and 11 references, 6 of which are Soviet. 

ASSOCIATION: Moskovskiy institut tonkoy khimicheskoy tekhnologii im. M. V.
Lomonosova
(Moscow Institute of Fine Chemical Technology imeni M. V. Lomonosov)

SUBMITTED: November 5, 1958

Card 2/2

FERNOVSKIY, M. F. BC		17-4	
<p>Inheritance of nicotine and anabasine in interspecific hybrids of <i>Nicotiana glauca</i>, Gr. M. F. FERNOVSKIY, M. I. CHMURA, and N. I. ZUKOV (Compt. rend. Acad. Sci. U.R.S.S., 1937, 17, 43—45).—Hybrids of <i>N. tabacum</i> (0.85% nicotine) with <i>N. glauca</i> (0.6% anabasine) contain no nicotine but 0.55—1.28% of anabasine. Hybrids of <i>N. glauca</i> × <i>N. rustica</i> and triple hybrids of these and <i>N. sylvestris</i> show a general tendency towards an increase in anabasine at the expense of nicotine in comparison with the parent plants. A. O. P.</p>			
ASS-5LA METALLURGICAL LITERATURE CLASSIFICATION			
FROM SYNONYM		FROM SYNONYM	
SECONDARY ONLY ONE		SECONDARY ONLY ONE	
CATEGORY		CATEGORY	
CATEGORY		CATEGORY	

TERNOVSKIY

M-F

AM

TERNOVSKY (M. F.) & KHUDYNA (I. P.). Отношение гибридов *Nicotiana glutinosa* L. x *N. tabacum* L. к. обычной Табачной мозаике. Предварительное сообщение. [Reaction of *Nicotiana glutinosa* L. x *N. tabacum* L. hybrids to ordinary Tobacco mosaic. Preliminary report.]-Вестник, научно-исп. Пром. Табачн. Мазорчи. Пром. им. А. И. Миколы (ВУТИМ) [The A.I. Mikolay prin-Soviet sci. Res. Inst. Tob. and Indian Tob. Ind. (VITIM)], Krasnodar, Publ. 135, pp. 69-70, 1938.

The annual losses from tobacco mosaic in the Azoff-Black Sea region are stated to amount to 3 to 5 per cent. In resistance trials in 1935 and 1936 no variety of *Nicotiana tabacum* was found to be immune from mosaic, but *N. glutinosa* developed necrotic lesions only in the region of inoculation (localized infection) and its leaves remained healthy and did not contain the virus. In breeding against tobacco mosaic [R.I.M., xvii, p. 417 and next abstract] Ternovsky succeeded

ASB.SLA METALLURGICAL LITERATURE CLASSIFICATION

6500 65100000

6500 65100000

6500 65100000

6500 65100000

6500 65100000

6500 65100000

in 1932 in obtaining amphidiploids from the cross *N. glutinosa* × *N. tabacum* with 2 genomes of *N. glutinosa* (24 chromosomes) and 2 genomes of *N. tabacum* (48 chromosomes). Back-crossing to *N. glutinosa* produced a sesquidiploid with 2 genomes of *N. glutinosa* and 1 genome of *N. tabacum*; back-crossing to *N. tabacum* produced a sesquidiploid with 2 genomes of *N. tabacum* and 1 genome of *N. glutinosa*.

Inoculation experiments showed that in the first generation hybrids both sesquidiploids exhibited only local infection; the second generation from the sesquidiploid with 2 genomes of *N. tabacum* segregated in the number of chromosomes and in resistance to mosaic; the majority of the plants showing general infection; in the third generation from the sesquidiploid with 2 genomes of *N. tabacum*, 10 out of 19 examined families were found to contain resistant plants, one of these lines showing 50 per cent. plants with local infection.

FERNOVSKIY, M. F.		PROCESSING AND PROPERTIES INDEX																																																																																																																																																																																																									
Am		<p>ТЕРНОВСКИЙ (М. Ф.). Наследственность локализации мозаики у гибридов <i>Nicotiana glutinosa</i> L. x <i>N. tabacum</i> L. — [Inheritance of mosaic localization in <i>Nicotiana glutinosa</i> L. x <i>N. tabacum</i> L. hybrids.] — Вестник науки и техники. Рязань. Табачн. Магистрат. Пром. им. А. Н. Мухоморова (ВНТМ). [The A.I. Mikhomorov pan-Soviet sci. Res. Inst. Tob. and Indian Tob. Ind. (VITIM)]. Krasnodar, Publ. 133, pp. 71-74, 1938. (English summary.)</p> <p>Two years' study on the susceptibility of <i>Nicotiana glutinosa</i> x <i>N. tabacum</i> hybrids to mosaic (see preceding abstract) showed that the</p>																																																																																																																																																																																																									
ASB-55A METALLURGICAL LITERATURE CLASSIFICATION		FROM SOURCE																																																																																																																																																																																																									
<table border="1"> <tr> <td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td><td>11</td><td>12</td><td>13</td><td>14</td><td>15</td><td>16</td><td>17</td><td>18</td><td>19</td><td>20</td><td>21</td><td>22</td><td>23</td><td>24</td><td>25</td><td>26</td><td>27</td><td>28</td><td>29</td><td>30</td><td>31</td><td>32</td><td>33</td><td>34</td><td>35</td><td>36</td><td>37</td><td>38</td><td>39</td><td>40</td><td>41</td><td>42</td><td>43</td><td>44</td><td>45</td><td>46</td><td>47</td><td>48</td><td>49</td><td>50</td><td>51</td><td>52</td><td>53</td><td>54</td><td>55</td><td>56</td><td>57</td><td>58</td><td>59</td><td>60</td><td>61</td><td>62</td><td>63</td><td>64</td><td>65</td><td>66</td><td>67</td><td>68</td><td>69</td><td>70</td><td>71</td><td>72</td><td>73</td><td>74</td><td>75</td><td>76</td><td>77</td><td>78</td><td>79</td><td>80</td><td>81</td><td>82</td><td>83</td><td>84</td><td>85</td><td>86</td><td>87</td><td>88</td><td>89</td><td>90</td><td>91</td><td>92</td><td>93</td><td>94</td><td>95</td><td>96</td><td>97</td><td>98</td><td>99</td><td>100</td> </tr> </table>		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100	<table border="1"> <tr> <td>1</td><td>2</td><td>3</td><td>4</td><td>5</td><td>6</td><td>7</td><td>8</td><td>9</td><td>10</td><td>11</td><td>12</td><td>13</td><td>14</td><td>15</td><td>16</td><td>17</td><td>18</td><td>19</td><td>20</td><td>21</td><td>22</td><td>23</td><td>24</td><td>25</td><td>26</td><td>27</td><td>28</td><td>29</td><td>30</td><td>31</td><td>32</td><td>33</td><td>34</td><td>35</td><td>36</td><td>37</td><td>38</td><td>39</td><td>40</td><td>41</td><td>42</td><td>43</td><td>44</td><td>45</td><td>46</td><td>47</td><td>48</td><td>49</td><td>50</td><td>51</td><td>52</td><td>53</td><td>54</td><td>55</td><td>56</td><td>57</td><td>58</td><td>59</td><td>60</td><td>61</td><td>62</td><td>63</td><td>64</td><td>65</td><td>66</td><td>67</td><td>68</td><td>69</td><td>70</td><td>71</td><td>72</td><td>73</td><td>74</td><td>75</td><td>76</td><td>77</td><td>78</td><td>79</td><td>80</td><td>81</td><td>82</td><td>83</td><td>84</td><td>85</td><td>86</td><td>87</td><td>88</td><td>89</td><td>90</td><td>91</td><td>92</td><td>93</td><td>94</td><td>95</td><td>96</td><td>97</td><td>98</td><td>99</td><td>100</td> </tr> </table>		1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100																																																																																																								
1	2	3	4	5	6	7	8	9	10	11	12	13	14	15	16	17	18	19	20	21	22	23	24	25	26	27	28	29	30	31	32	33	34	35	36	37	38	39	40	41	42	43	44	45	46	47	48	49	50	51	52	53	54	55	56	57	58	59	60	61	62	63	64	65	66	67	68	69	70	71	72	73	74	75	76	77	78	79	80	81	82	83	84	85	86	87	88	89	90	91	92	93	94	95	96	97	98	99	100																																																																																																								

R2 AM TERNOVSKIY, M.F.

ТЕРНОВСКИЙ (М. Ф.). Методика селекции невосприимчивых сортов Табака к табачной мозаике и мучнистой росе. [Methods of breeding Tobacco varieties resistant to Tobacco mosaic and powdery mildew.]—Всесоюз. научисслед. Инст. Табачн. Материч. Прав. им. А. И. Микояна (ВНТИМ). [The A. I. Mikoyan Inst.-Soviet sci. Res. Inst. Tob. and Indian Tob. Ind. (VITIM)], Krasnodar, Publ. 143, pp. 126-141, 1941. [Received December, 1944. English summary.]

The author outlines the methods of preparing the virus solution for artificial infection, the mode and time of infection, and the manner of calculating the data used in his breeding work with hybrids of *Nicotiana glutinosa* and *N. tomentosiformis*, which showed a promising degree of resistance to tobacco mosaic and powdery mildew (*Erysiphe cichoracearum*: R.A.M., xx, p. 89).

TERNOVSKIY, M.F.

GERS L No. 43

Ternovskiy, M.F. (A.I. Mikoyan All-Union Scientific Research Institute of Tobacco), changes of resistance to tobacco mosaic as a result of vegetative hybridization, 517-20

Akademiya Nauk S.S.S.R. Doklady Vol. 79 No. 3, 1951

B-5

USSR / General Biology. Genetics

Abs Jour : Ref Zhur - Biol., No 11, 1958, No 47621

Author : Ternovskiy, M. F.

Inst : Not given

Title : The Introduction of Resistant Varieties of Tobacco.

Orig Pub : Michurinsk Sb Krasnodar, 'Sov Kuban', 1957, 158-168.

Abstract : The wild-growing variety *Nicotiana glutinosa* which is resistant to tobacco mosaic and to *Erysiphe graminis* was crossed with cultivated tobacco varieties late in the fall when a lowering of the temperature had already set in. The F_1 showed 1-2% amphidiploids. The latter were found to be resistant to both diseases but the quality of the production obtained from them was low. The amphidiploids were crossed with the better quality cultured varieties. The subsequent generations selected for high agrotechnical indices were repeatedly crossed with cultivated varieties. Resistance was

Card 1/2

USSR / General Biology. Genetics.

B-5

Abs Jour : Raf Zhur - Biol., No 11, 1958, No 47621

Abstract : always found to be dominant over susceptibility. The immunity was observed to be operative at all stages of development of the plants, resulting in the development of varieties resistant to both tobacco mosaic and *Erysiphe graminis*. The resistant new variety Trapezond 161 in epidemic years out-yields the unresistant former standard Trapezond 1272 by 20-50%. The new resistant variety Alma-Ata Dyubek 7 outyields the previous standard Dyubek 44 by 33% on the average and also yields a better quality tobacco. P. M. Grushevaya has obtained by crossing cultivated tobacco varieties with *N. glutinosa*; the new variety is resistant to both tobacco mosaic and to black root rot.

Card 2/2

COUNTRY : USSR
CATEGORY :

ABST. JOUR. : RZBiol., No. 19, 1958, No. 87182

AUTHOR : Ternovskiy, M. F.

INST. :
TITLE : The Utilization of Wild Species of Tobacco
in Selection for Disease Resistance.

ORIG. PUB. : Vestn. s.-Kh. nauki, 1958, No 1, 137-139

ABSTRACT : Tobacco of the Dyubek 44 variety was crossed with a wild species *Nicotiana glutinosa*, which has the capability of localizing the virus of tobacco mosaic by forming necrotic spots on the leaves. There were obtained a small number of constant-intermediate hybrid-amphidiploids (with a doubled set of chromosomes) which were not susceptible to tobacco mosaic but exhibited a number of undesirable characteristics. For their elimination use was made of repeated impregnation of fertile hybrids with tobacco, and continuous selection for immunity, plant vigor, degustatory characteristics and leaf color. In this manner were obtained, for the first time, tobacco

CARD:1/3

Country : USSR
CATEGORY :

M-7

ABR. JOUR. : RZBiol., No. 19, 1958⁸, No. 87182

AUTHOR :
INST. :
TITLE :

ORIG. PUB. :

ABSTRACT : varieties that are immune to tobacco mosaic. The same fertile hybrids between tobacco and *N. glutinosa* were used to produce varieties that are immune to parasitic fungus *Erysiphe graminis*. As a result, completely immune tobacco varieties were developed: Dyubek 566, Damsun 47/10 Alma-Atinskii 315, Amerikan 287, Trapezond 141, Trapezond Talasskiy, Immunnyy 3000. Some of them have already been put in production. The development of multiple-immunity tobacco varieties is an important achievement of Soviet science. Resistance to disease of the wild species *N. glauca* could not be put to use, because of considerable difficulty of crossing it with tobacco and the sterility of first generation hybrids. By crossing the resistant wild

CARD: 2/3

COUNTRY : USSR
CATEGORY :

M-7

ABR. JOUR. : RZBiol., No. ⁸19, 1959, No. 87182

AUTHOR :
INST. :
TITLE :

ORIG. PUB. : .

ABSTRACT : species *N. glauca* with tobacco and *N. rustica*, fertile hybrids were obtained, some of which can serve as starting material for the development of varieties of *N. rustica* that are resistant to top chlorosis (virus disease of *N. rustica*). Hybridization between species can also be successfully utilized for the production of forms of tobacco and *N. rustica* which are resistant to other diseases and infection. The work was carried out by the All-Union Scientific Research Institute of Tobacco and *N. rustica*. -- L. A. Lomakina.

CARD: 3/3

611

TERNOVSKIY, M.F.; TEREENT'YEVA, A.I.

Role of grafting in increasing the crossability of *Nicotiana* species. Dokl.AN SSSR 132 no.4:932-935 Jo '60. (MIRA 13:5)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut tabaka i makhorki, g. Krasnodar. Predstavleno akademikom N.V.TSitsinyam.
(Tobacco breeding) (Grafting)

ANTONENKO, G.P., agronom; KHOKHRYAKOV, M.K., prof.; TERNOVSKIY, M.P., prof.

Peronospora (downy mildew) infection of tobacco in Czechoslovakia.
Zashch. rast. ot vred. i bol. 6 no.5:53-54 My '61. (MIRA 15:6)
(Czechoslovakia—Tobacco blue mold)

TERNOVSKIY, M. F., prof.

Tobacco varieties unsusceptible to diseases. Zashch. rast.
ot vred. i bol. 6 no.6:26-27 Je '61. (MIRA 16:4)

1. Vsesoyuznyy institut tabaka i makhorki, Krasnodar.

(Tobacco—Disease and pest resistance)

TERNOVSKIY, M.F.

Polyploidy in the genus Nicotiana. Trudy MOIP.Otd.biol. 5:230-
237 '62. (MIRA 16:5)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut tabaka i makhorki,
Krasnodar. (TOBACCO BREEDING) (POLYPLOIDY)

TERNOVSKIY, M.F.; DASHKEYEVA, K.N.

Responses of the species *Nicotiana* to infection with false tobacco mildew. Dokl. AN SSSR 150 no.4:931-933 Je '63.
(MIRA 16:6)

1. Institut fiziologii i biokhimii rasteniy AN Moldavskoy SSR. Predstavleno akademikom N.V. TSitsinyam.
(Tobacco blue mold)
(Tobacco—Diseases and pests)

TERNOVSKIY, M.F., doktor sel'skokhoz.nauk; DASHKEYEVA, K.N., kand.biolog.
nauk

Estimating the resistance of tobacco to downy mildew. Zashch. rast.
ot vred. i bol. 9 no.1:20-23 '64. (MIRA 17:4)

1. Vsesoyuznyy nauchno-issledovatel'skiy institut tabaka i makhorki
imeni A.I.Mikoyana, Krasnodar, i Institut fiziologii i biokhimii
rasteniy AN Moldavskoy SSR, Kishinev.

TERNOVSKIY, M.F.; DASHKEYEVA, K.N.

Wild species of tobacco as a source for developing varieties
resistant to downy mildew. Izv.AN Mold.SSR no.4:63-74 '63.
(MIRA 18:1)

TERNOVSKIY, N.S.

Operation of the Lipetsk Alcohol Plant with continuous cooking.
Spir. prom. 24 no. 4:21-23 '58. (MIRA 11:7)
(Alcohol)

TERNOVSKIY, N.S.

Prague symposium on the subject: Continuous cultivation of
micro-organisms. Spirt. prom. 24 no.6:47-48 '58. (IRA 11:10)
(Prague--Micro-organisms--Congresses)

TERNOVSKIY, N.S.

Session of the Scientific Council of the All-Union Scientific
Research Institute of the Food Engineering Equipment. Spirt.
prom. 26 no.1:43-45 '60. (MIRA 13:6)
(Food industry--Equipment and supplies)

TERNOVSKIY, N.S.; GRYAZNOV, V.P.

Beer rectification columns used in France. Spirt.prom.
26 no.4:27-31 '60. (MIRA 13:8)
(France--Distilling industries--Equipment and supplies)

GRYAZNOV, V.P.; BOGDANOV, Yu.P.; RZHECHITSKAYA, G.V.; TERNOVSKIY, N.S.;
GRACHEV, B.K. [deceased] MERKIN, V.G.; POLEVAYA, K.G.;
AKIMENKO, I.S.

Double-flow beer rectification apparatus. Spirt. prom. 28
no.7:35-37 '62. (MIRA 17:2)

1. Tsentral'nyy nauchno-issledovatel'skiy institut spirtovoy i
likero-vodochnoy promyshlennosti (for Gryaznov, Bogdanov,
Rzhechitskaya, Ternovskiy). 2. Lipetskiy spirtovoy zavod (for
Grachev, Merkin, Poleyaya, Akimenko).

DENSHCHIKOV, Mikhail Tikhonovich, kand.tekhn.nauk; SILIN, P.M., prof., red.; VESELOV, I.Ya., prof., red.; SMIRNOV, V.A., prof., red.; RZHEKHIN, V.P., red.; LEEDEV, P.P., red.; KOVALENKO, Yu.T., red.; KUPCHINSKIY, P.D., red.; BENIN, G.S., red.; P'YANKOV, A.G., red.; SHNAYDMAN, L.O., red.; MOREV, N.Ye., red.; SHMAIN, M.M., red.; BULGAKOV, N.I., red.; MAYOROV, V.S., red.; TERNOVSKIY, N.S., red.; RAZUVAYEV, N.I., red.; OGORODNIKOV, S.T., red.; BURMAN, M.Ye., red.; KHOLOSTOV, V.A., red.; NAMESTNIKOV, A.F., red.; NASAKIN, T.N., red.; KOVALEVSKAYA, A.I., red.; KISINA, Ye.I., tekhn. red.

[Wastes from the food industry and their utilization] Otkhody pishchevoi promyshlennosti i ikh ispol'zovanie. Izd. 2., dop. i perer. Moskva, Pishchepromizdat, 1963. 615 p. (MIRA 16:6)
(Food industry--By-products)

TERNOVSKIY, N. S.; LOSIK, L. I.

Development and manufacture of the new type of technological equipment for the distilling industries. Spirt. prom. 29 no.3: 22-24 '63. (MIRA 16:4)

1. Gosudarstvennyy komitet po avtomatizatsii i mashinostroyeniyu pri Gosplane SSSR (for Ternovskiy). 2. Gosudarstvennyy institut po proyektirovaniyu predpriyatiy spirtovoy i vinodel'cheskoy promyshlennosti (for Losik).

(Distilling industries—Equipment and supplies)

TERNOVSKIY, N.S.; GRYAZNOV, V.P.

Experience in the thermal processing of alcohol in the Michurinsk
and Lipetsk distilleries. Fern. i spirt. prom. 31 no.2:36-38 '65.
(MIRA 18:6)

1. Gosudarstvennyy komitet po pishchevoy promyshlennosti (for
Ternovskiy). 2. Vsesoyuznyy nauchno-issledovatel'skiy institut
fermentnoy i spirtovoy promyshlennosti (for Gryaznov).

TERNOVSKIY, P. (Alma-Ata)

Improve the quality of photographic information. Sov.foto. 23
no.2:45 F '63. (MIRA 16:4)

(Photography)

TERNOVSKIY, S.D.

DECEASED
c1960

1962/3

SEE ILC

PEDIATRICS

TERNOVSKIY, V.A., dotsent, kandidat tekhnicheskikh nauk, redaktor; MATVEYEVA, E.N., tekhnicheskiiy redaktor.

[Motion locomotives on curved sections of track. Stability in the movement of locomotives] Vpisyvanie lokomotivov v krivye uchastki zheleznodorozhnogo puti. Ustoichivost' dvizhenia lokomotivov. Moskva, Gos. nauchno-tekhn. izd-vo mashinostroit. lit-ry . Pt. 1. 1954. 112 p. (MIRA 8:1)

1. Russia (1923- U.S.S.R.) Ministerstvo vysshego obrazovaniia. (Railroads--Curves and turnouts) (Locomotives--Design)

TERNOVSKIY, V.A., kand.tekhn.nauk; STRAMOUS, M.F., kand.tekhn.nauk

"Track machinery and mechanisms" by N.N.Gulenko, V.E.Gora.
Reviewed by V.A.Ternovskii, M.F.Stramous. Put' i put.khoz. 6
no.3:46 Mr '62. (MIRA 15:3)
(Railroads--Equipment and supplies)
(Gulenko, N.N.) (Gora, V.E.)

DERBAREMDI~~ER~~, M.I.; SEREBRENNIKOVA, K.L.; TERNOVSKIY, V.A.; Prinsipal
uchastiye: SHAROV, P.M.; NOVIKOV, L.Z.; LUR'YE, E.Y.; PIS'MEN,
M.K.; KARABIN, A.I. [deceased]; KOSTIN, L.I.; FROLOV, V.P.;
MEDVEDEV, F.V.; GELIMKHANOV, S.G.; BONDAR', V.G.; TIMOFEYEV,
P.I.; MININA, L.V.; AREEKOV, F.F.; NIKOLAYEV, N.I.; YAROSLAV,
T.Ye.; NUDEL'MAN, V.G.

Gasification of mazut under pressure in a steam-oxygen blast.
Gaz. prom. 9 no.11:49-50 '64. (MIRA 17:12)

TERNOVSKIY V. N.

A bibliography of dissertations of the Faculty of Medicine of Moscow University.

1949 63 p.

DAFM

1. Dissertations, Academic - Russia - Bibl.
2. Medicine - Bibl.
- I. Moscow. Universitet. Meditskinskii fakul'tet.

Problems in morphology; collection of scientific works Moskva, Izd-vo Akademii med.
nauk SSSR, 1949. 108 p. (52-37722)

QM23.T268

TERNOVSKIY, V. N.

21996 TERNOVSKIY, V. N. i MURAT, V. N. O metodike izucheniya sosudistyykh taseynov krovesnab- zheniya chreanov. Vsb: Voprasy morfologii, M., 1949, s. 24-28.

SO: Letopis' Zhurnal'nykh Statey, No. 29, Mskva, 1949.

1. TERNOVSKIY, V. N.

2. USSR (600)

4. Avicenna, 980? - 1037

7. Avicenna and his great accomplishment, the Canon of Medicine.
Arkhir. anat. gist. i embr. 29. No. 5. 1952.

9. Monthly List of Russian Accessions, Library of Congress, February 1953. Unclassified.

TERNOVSKIY, V.N.

[Renaissance medicine and its study in the Soviet Union.]
Meditsina epokhi vrozozhdenia i ee isuchenie v Sovetskom
Soiuze. Moskva, Medgiz, 1954. 22 p. ports. (MLRA 9:1)
(MEDICINE--HISTORY)

TERNOVSKIY V.N.
VESALIUS, Andreas; **TERNOVSKIY**, V.N., redaktor, [translator]; **SHESTAKOV**, S.P., [translator]; **PAVLOV**, I.P., akademik; **PETROVSKIY**, I.G., akademik, redaktor; **BYKOV**, K.M., akademik, redaktor; **KAZANSKIY**, B.A., akademik, redaktor; **OPARIN**, A.I., akademik, redaktor; **SHMIDT**, O.Yu., akademik, redaktor; **ANDREYEV**, N.N., akademik, redaktor; **KOSHTOYANTS**, Kh.S., redaktor; **SAMARIN**, A.M., redaktor; **MAKSIMOV**, A.A., redaktor; **SHCHERBAKOV**, D.I., akademik, redaktor; **YUDIN**, P.F., akademik, redaktor; **LEBEDEV**, D.M., doktor geograficheskikh nauk, redaktor; **FIGUROVSKIY**, N.A., doktor khimicheskikh nauk, redaktor; **KUZNETSOV**, I.V., kandidat filosofskikh nauk, redaktor; **OZNOBISHIN**, D.V., kandidat istoricheskikh nauk, redaktor; **SHIDLOVSKAYA**, O.G., redaktor; **RUDNEVA**, O.A., redaktor; **KISELEVA**, A.A., tekhnicheskii redaktor.

[Structure of the human body; in 7 books] O stroenii chelovecheskogo tela; v semi knigakh. Perevod s latinskogo V.N.Ternovskogo i S.P.Shestakova. Red. V.N.Ternovskogo. Posleslovie I.P.Pavlova. [Moskva] Izd-vo Akademii nauk SSSR. Vol. 2. 1954. 960 p. (MIRA 7:11)

1. Deystvitel'nyy chlen Akademii meditsinskikh nauk SSSR (for Ternovskiy)
2. Chlen-korrespondent Akademii nauk SSSR (for Shestakov, Koshtoyants, Samarin, Maksimov)
 (Anatomy, Human--Early works to 1800)

TERNOVSKIY, V.N.
TERNOVSKIY, V.N.

Aleksai Nikolaevich Gennadiev; 1892-1956. Arkh.anat.gist. 1 embr. 34
no.5:126 S-O '57. (MIRA 11:1)

(GENNADIEV, ALEKSEI NIKOLAEVICH, 1892-1956)

TERNOVSKIY, V.N., professor (Moskva)

Portrait of Avicenna. Klin. med. 35 no.1:120-121 Ja '57
(MIRA 10:4)

1. Deystvitel'nyy chlen Akademii meditsinskikh nauk SSSR.

(FAMOUS PEOPLE

Avicenna, portrait according to photodocumentation
of skull)

TERNOVSKIY, V.N.

Ibn Sina and the reproduction of his portrait. Vest.oft. 70 no.6:42
N-D '57. (MIRA 11:1)

1. Deystvitel'nyy chlen AMN SSSR.
(AVICENNA, 9807-1037)

ABU ALI IBN SINA (Avicenna); TERNOVSKIY, V.N., pref., otv.red.; ARENDS, A.K., kand.filol.nauk, otv.red.; SHIPUKHIN, A.Ya., red.isd-va; GOR'KOVAYA, Z.P., tekhn.red.

[Canon of medicine] Kanon vrachebnoi nauki. Tashkent, Izd-vo Akad.nauk Uzbekskoi SSR. Book 3, vol.1. 1958. 793 p.
(MIRA 12:3)

1. Deystvitel'nyy chlen Akademii Meditsinskikh nauk SSSR (for Ternovskiy).

(MEDICINE, ARABIC)

TERNOVSKIY, V.N., prof.

Sixteenth International Congress of Medical Historians. Vest.
AMN SSSR 14 no.8:68-71 '59. (MIRA 12:11)

1. Deystvitel'nyy chlen AMN SSSR.
(MEDICINE)

ABU ALI IBN SINA (AVICENNA) [deceased]; SAL'YE, M.A., kand.filol.nauk, starshiy nauchnyy sotrudnik [translator]; TERNOVSKIY, V.N., prof., akademik, otv.red.; PETROV, B.D., kand.med.nauk, red.; ASKAROV, A.A., red.; KARIMOV, U.I., kand.filol.nauk, red.; AZIMDZHANOVA, S.A., kand.istor.nauk, red.; ARENDS, A.K., kand.filol.nauk, red.; DZHUMAYEV, V.K., kand.med.nauk; RASULEV, A., starshiy.nauchnyy sotrudnik; MIL'MAN, Z.A., red.; GOR'KOVAYA, Z.P., tekhn.red.

[Canon of medical science] Kanon vrachebnoi nauki. Tashkent, Izd-vo Akad.nauk Uzbekskoi SSR. Book 4. 1960. 767 p.

(MIRA 13:12)

1. Institut vostokovedeniya AN UzSSR (for Sal'ye).
2. Akademiya meditsinskikh nauk SSSR (for Ternovskiy).
3. Zaveduyushchiy kafedroy istorii meditsiny Moskovskogo meditsinskogo instituta (for Petrov).
4. Zaveduyushchiy laboratoriyey Instituta krayevoy meditsiny, chlen-korrespondent AN UzSSR (for Askarov).

(MEDICINE, ARABIC)

ABU ALI IBN SINA (AVICENNA) [deceased]; KARIMOV, U.I., kand.filolog. nauk [translator]; TERNOVSKIY, V.M., prof., akademik, otv.red.; ARENDS, A.K., kand.filolog.nauk, otv.red.; PETROV, B.D., kand.med. nauk, red.; AZIMDZHANOVA, S.A., kand.istor.nauk, red.; ASKAROV, A.A., red.; DZHUMAYEV, V.K., kand.med.nauk, red.; KARASIK, V.M., red.; RASULEV, A., starshiy nauchnyy sotrudnik, red.; MIL'MAN, Z.A., red.; BABAKHANOVA, A.G., tekhn.red.

[Canon of medical science] Kanon vrachebnoi nauki. Tashkent, Izd-vo Akad.nauk Uzbekskoi SSR. Book 5. 1960. 329 p.

(MIRA 13:12)

1. Zaveduyushchiy otdelom nauchnogo opisaniya i katalogizatsii rukopisey Instituta vostokovedeniya Akademii nauk UzSSR (for Karimov).
2. Akademiya meditsinskikh nauk SSSR (for Ternovskiy).
3. Zaveduyushchiy otdelom izucheniya i publikatsii rukopisnykh pamyatnikov Instituta vostokovedeniya AN UzSSR (for Arends).
4. Zaveduyushchiy kafedroy istorii meditsiny Moskovskogo meditsinskogo instituta (for Petrov).
5. Chlen-korrespondent AN UzSSR (for Askarov).
6. Deyatel'nyy chlen Akademii meditsinskikh nauk SSSR (for Karasik).
7. Institut vostokovedeniya AN UzSSR (for Rasulev).

(MEDICINE, ARABIC)

TERNOVSKIY, V.N., prof.

Tasks of the Academy of Medical Sciences of the U.S.S.R. in
publishing medical classics. Vest.AMN SSSR 15 no.1:68-70 '60.
(MIRA 13:6)

1. Deyatvitel'nyy chlen AMN SSSR.
(MEDICINE)

TERNOVSKIY, V.N. (Moskva, Novoslobodskaya ul., 57/65, kv. 34)

Nikolai Bidloo; on the 225th anniversary of his death. Arkh.anat.
gist.i embr. 39 no.7:87-90 J1 '60. (MIRA 14:5)
(BIDLOO, NIKOLAI, d.1735)

TERNOVSKIY, V.N.

The 17th International Congress of Medical Historians. Vent. AMN
SSSR 16 no.8:84-87 '61. (MIRA 14;12)
(MEDICINE... CONGRESSES)

TERNOVSKIY, V.N.

Complete edition of Avicenna's "Canon of medical science." Reviewed
by V.N.Ternovskii. Sov.zdrav. 20 no.4:69-73 '61. (MIRA 14:5)

1. Otvetstvennyy redaktor "Kanona" deystvitel'nyy chlen Akademii
meditsinskikh nauk SSSR.
(AVICENNA, 980? - 1037?)

TERNOVSKIY, V.P., zasluzhennyy vrach RSFSR (Penza)

Otorhinolaryngological services in Penza Province and prospects
for their improvement. Vest.otorin. 21 no.4:79-81 J1-Ag
'59. (MIRA 12:10)
(OTORHINOLARYNGOLOGY hosp. & clinics)

TERNOV-TROP, L.K., izn.; ROYZ, I.A., izn.; SOGOLOV, I.I., izn.

Hammer mill clutches. Energetik 12 no.7:17-20 J1 '64.
(MIRA 17:9)

SHALIMOVA, K.V.; VERDANYAN, V.I.; TERNOVYKH, Yu.P.

Electric and optical properties of thin layers of germanium and
silicon. Nauch.dokl.vys.shkoly; radiotekh.i elektron. no.4:232-238
'58. (MIRA 12:6)

1. Kafedra poluprovodnikovyykh priborov Moskovskogo energeticheskogo
instituta.

(Germanium) (Silicon)

TERNOVYY, M. [Ternovyi, M.]

Circular mechanized seine feeder. Sil'.bud. 13 no.10:6-7 0 '63.
(MIRA 17:3)

1. Glavnyy inzh.-stroitel' Khmel'nitskogo kol'khozno-sovkhoznogo
proizvodstvennogo upravleniya Vinnitskoy oblasti.

TERNOVYY, Yu. [Ternoyi, IU.] (Moskva)

About the miracle clock and the "Grand Prix." Znan. ta pratsia
no. 1:11-12 Ja '61. (MIRA 14:4)
(Moscow—Vocational Education—Exhibitions)

TERNOVYY, Yu. [Ternovyl, IU.]

Three equators of the Dnieper Valley Little Railroad. Znan.ta
pratsia no.8:12 Ag '62. (MIRA 15:12)
(Dnieper Valley—Railroads)

SZUCS, Miklos, dr., a kémiai tudományok kandidátusa; SZASZ Frigyes;
BUNYITAI, Janos, dr.; BANAS Jozsef; ZACHETSKY, Ferenc;
GULDEN, Otto; TERNYAK, Beno; BENKOVIC, Jozsef; DIOSZEGHY,
Daniel, prof., dr.; TOTH, Istvan.

Questions of vocational education pertaining to the gas in-
dustry. Energia es atom 17 no.1:7-11 Ja'64.

1. Fovarosi Gazmuvek (for Szucs); 2. Orszagos Koolaj- es
Gazipari Troszt (for Szasz and Bunyitai). 3. Miskolci Ne-
hezipari Muszaki Egyetem (for Dioszeghy). 4. Veszpremi
Vegyipari Muszaki Egyetem (for Toth).

KISS, Lorant, okleveles gepeszmernok; CSERNAVOLGYI, Laszlo; HAJDU, Istvan;
BENKOVICS, Jozsef; TERNYAK, Beno; SOSKUTI, Andras; TOROK, Mihaly, dr.;
SZASZ Frigyes; GATI, Geza; KOVACS, Lajos; DEHENES, Zoltan; MAGYAR, ~~Laszlo~~,
Laszlo; KOVACS, Gyula; AUERSWALD, Janos; SOS, Janos; DIOSZEGHY, Daniel,
prof.

Manufacture and use of gas appliances. Energia es atom 17 no.1:
30-35 Ja'64.

1. Lampagyar (for Kiss).
2. Vegyterv (for Csernavolgyi).
3. Orszagos Koolaj- es Gazipari Troszt (for Hajdu, Szasz, Auerswald).
4. Pecsai Gazszolgaltato Vallalat (for Benkovics).
5. Asvanyolajforgalmi Vallalat (for Ternyak, Soskuti).
6. Epitesugyi Miniszterium Iparterv Musze'ki Osztaly (for Torok).
7. Orszagos Villamosenergia Felugyelet (for Gati).
8. Epitesugyi Miniszterium (for Lajos Kovacs).
9. Gazkeszulekgyarto Vallalat (for Dehenes).
10. Epitestudomayi Intezet (for Gyula Kovacs).

CSIBA, Lajos; VASARHELYI, Istvan; RADETSKY, Jeno; PATKAI, Imre, dr.; TERNYAK,
Jeno; SCHAFER, Lajos

Data on the birds of prey. Aquila 69/70:258 '62-'63 [publ. '64].

JAKAB, Andras; TERNYAK, Jeno; PATKAI, Imre; KALOCZY, Lajos; CSIBA, Lajos

Swan occurrences. Aquila 69/70:257 '62-'63 [publ. '64].

SCHMIDT, Egon; STERBETZ, Istvan; GYERESSY, Antal; SCHAFER, Lajos; TERNYAK, Jeno;
MATE, Laszlo; GEREBY, Gyorgy; BERETZK, Peter, dr.

Data on the avifauna of the region between the Danube and the
Tisza. Aquila 69/70:258-260 '62-'63 [publ. '64].

MASLOV, V.A., inzh.; TERNYUK, M.I., inzh.; GERMAN, V.T., inzh.

Effect of deformation on the resistance to corrosion of type
18-8 steels. Khim.mashinostr. no.1;28-30 Ja-F. '64. (MIRA 17:4)

SOV/20-122-3-34/57

AUTHORS: Gindin, L. M., Bobikov, P. I., Kouba, E. F., Kopp, I. F., Rozen, A. M., Ter-Oganesov, N. A., Zagarskaya, N. I.


TITLE: Separation of Metals by the Exchange-Extraction Method
(Razdeleniye metallov metodom obmennoy ekstraktsii)

PERIODICAL: Doklady Akademii nauk SSSR, 1958, Vol 122, Nr 3, pp 443-447
(USSR)

ABSTRACT: An extraction in connection with an exchange reaction between metals is a very productive method of separation if these metals are in different phases: in an organic phase as salts and aliphatic acids and in an aqueous phase as salts of mineral acids (Ref 1). For this purpose saturated aliphatic acids with 5 and more carbon atoms were used. They fulfill a double function: a) they take part in the formation of the corresponding metallic salts (soaps), and b) they serve as solvents for these soaps being formed. Aliphatic acids are used most properly as solutions in an inactive solvent with a low specific weight. Directions for the preparation of such solutions are mentioned. The exchange reaction between the metals as mentioned earlier can be expressed by the following equation:

Card 1/3

Separation of Metals by the Exchange-Extraction Method SOV/20-122-3-34/57



^1M and ^2M denote the corresponding metals, R - the organic residue of the aliphatic acid $\text{C}_n\text{H}_{2n+1}\text{COO}'$, the indices org and aqu denote the organic and the aqueous phase. The equilibrium constant of the exchange reaction depends on the character of the exchanging metals, as was confirmed by the experiments. Metals with a small pH value ("acid" metals) mainly pass into the organic phase, metals with a high pH value, however, (more alkaline metals) into the aqueous phase. In many cases reaction (1) takes place almost completely (>99%), it may therefore be said that a metal is displaced from the organic phase by another metal. Separation of the metallic salts by means of the reaction mentioned in the title can be carried out from the aqueous as well as from the organic phase. In the first case (Fig 1) the aqueous phase which contains a mixture of salts of two metals is brought into contact with the organic phase in which a salt of an aliphatic acid of a stronger alkaline metal is contained. In the second case the organic phase which contains a mixture of salts of the aliphatic acids is brought into contact with the aqueous phase which contains a salt of a mineral acid of a

Card 2/3

Separation of Metals by the Exchange-Extraction Method SOV/20-122-3-24

weaker alkaline metal. Table 1 reveals the results of separation of metallic salts combined with sulfuric acid by means of the discussed method. As organic phase a solvent of industrial aliphatic acids of the fraction $C_7 - C_9$ (average molecular weight 141) in petroleum (400 g/liter) was used. Data on table 1 characterize a single exchange. By using an extraction column the degree of separation is considerably increased. If metals have similar properties reaction takes place incompletely. There are 2 figures, 1 table, and 1 reference, 1 of which is Soviet.

ASSOCIATION: Noril'skiy gorno-metallurgicheskiy kombinat im. A. P. Zavenyagina (Noril'sk Mining Metallurgy Kombinat imeni A. P. Zavenyagin)

PRESENTED: May 4, 1958, by S. I. Vol'fkovich, Member, Academy of Sciences. USSR

SUBMITTED: April 12, 1958

Card 3/3

GINDIN, L.M.; KOPP, I.F.; ROZEN, A.M.; BOBIKOV, P.I.; KOURA, E.F.;
TNR-OGANESOV, N.A.

Extraction equilibria for cobalt, nickel, and certain metals.
Zhur.neorg.khim. 5 no.1:149-159 Ja '60.

(MIRA 13:5)

1. Noril'skiy gornometallurgicheskiy kombinat im. A.P.
Zaveryagina, Opytno-issledovatel'skiy tsakh.
(Extraction (Chemistry)) (Metals)

TER-OGANESYAN, C.N., kand.med.nauk

Diagnostic importance of the pulsation of the abdominal aorta
in tuberculous mesoadenitis. Vop.rent.i onk. 6:333-337 '61.

(MIRA 16:2)

(ABDOMINAL AORTA) (PULSE) (LYMPHATICS—TUBERCULOSIS)

TER-OGANESYAN, I.M.

Measuring the resistance of the loop phase - zero. From.
energ. 16 no.7:17-21 J1 '61. (MIRA 15:1)
(Electric measurements)

TER-OGANESYAN, R.R.

Methods for determining the quality of insulation of underground
pipelines and organization of its systematic control. Trudy
Inst.energ.AN Gruz.SSR 16:193-204 '62. (MIRA 16:4)
(Electric lines--underground)

THE XIAN'SYAN, R.C.

Constant monitoring of the location of underground gas mains.
Gas. prom. V no. 112-47 '62 (MIRA 17.5)

TER-OGANESYAN, R.R.

Operation of current conductors with leakage and different
parameters in different sectors. Trudy Inst. energ. AN Gruz.
SSR 17:201-218 '63. (MIRA 17:7)

TER-OGANE\$OV, V. T.

Eclipses of the sun and the moon. Moskva, Gos. izd-vo kul'turno-prosvetitel'noi
lit-ry, 1952. 84 p. maps. (53-37193)

QB175.T45

TER-OGANESOV, Ya.G.; GVAYTA, T.I.; ROSHCIN, Yu.V.; ZUBOVA, V.I.

Method and equipment used in foreign countries in aerogeophysical
prospecting for uranium deposits. Atom. energ. Supplement no.6:146-160
'57. (MIRA 11:7)

(Aeronautics in surveying) (Prospecting--Geophysical methods)
(Uranium ores)

Ter-Oganesov, Ya. G.

AUTHORS: Ter-Oganesov, Ya. G., Gyayta, T. I., Roshchin, Yu. V. 89-1-21/29

TITLE: The Applications of Aeroradiometric Methods for the Detection of Workable Minerals (Primeneniye aeroradiometricheskikh metodov dlya poiskov razlichnykh poleznykh iskopaemykh)

PERIODICAL: Atomnaya Energiya, 1958, Vol. 4, Nr 1, pp. 102-102 (USSR)

ABSTRACT: With a few words the authors deal with the contents and the conclusions to be drawn from the following publications:
Aviation Week, 64, 2, p.74
Canad. Chem.Process, 37, 13, p. 66
Mines Mag., 46, 7, p. 31 (Kellog)
World Petrol, 23, 5, p. 109 (Lundberg)
Mining J., 234, p. 708 (1954)
Eng. and Mining J., 1954, Nr 7. p. 266
Photogram. Eng., 20, Nr 4, (1954)
There are 7 non-Slavic references.

AVAILABLE: Library of Congress

Card 1/1

SOV/11-59-6-15/15

30(7)

AUTHORS: Shcherbakov, D.I. and Ter-Oganesov, Ya.G.
TITLE: Scientific Ties with Belgian Scientists
PERIODICAL: Izvestiya Akademii nauk SSSR, Seriya geologicheskaya,
1959, Nr 6, pp 126-128 (USSR)
ABSTRACT: The authors describe their impression of the Brussels
World Exhibition and the contact they established
with Belgian scientists. Professor N.P. Yermakov and
geologist T.T. Matrenitskiy also took part in the
personal contacts with the Belgians.

Card 1/1

S/172/60/013/003-4/001/002
B002/B067

AUTHORS: Gvayta, T. I., Ter-Oganesov, Ya. G. (Moscow)

TITLE: Prospecting for Rare and Disperse Elements With Radioactive Methods a

PERIODICAL: Izvestiya Akademii nauk Armyanskoy SSR. Geologicheskkiye i geograficheskkiye nauki, 1960, Vol. 13, No. 3-4, pp. 57-63

TEXT: The present paper gives a general survey of the possibilities of prospecting for non-radioactive rare and disperse elements by radiometric methods. Many of these deposits also contain a certain amount of uranium and thorium which cause gamma anomalies in the deposits. Sometimes uranium and thorium minerals are paragenetically associated with the mineralization of rare and disperse elements, such as with beryllium and columbite - tantalite in pegmatites, with beryllium and wolframite in greisenizations; sometimes the uranium and thorium minerals themselves such as pyrochlore, lovtshorrite, fergusonite, monazite, xenotime, etc. are carriers of the rare elements. Besides, uranium and thorium may occur as isomorphous

Card 1/3

Prospecting For Rare and Disperse Elements
With Radioactive Methods

S/172/60/013/003-4/001/002
B002/B067

constituents in the gangue and in the accessories like in apatite, sphene, zircon, fluorite, and others. The magmatogenous deposits were divided according to the type of mineralization and the radiometric properties:

a) Premagmatic type: mineralization consists in finely distributed loparite, lovtshorrite, dysanlyte or knopite. Thorite, lovtshorrite etc. are radioactive minerals. The anomaly occurs locally and isometrically.

b) Post-magmatic type: The mineralization is related to late displacement processes, especially to albitization. The mineralized material consists of zircon (as cyrtolite and malacon) pyrochlore, polycrase or fergusonite. columbite. The radiometrical anomaly clearly follows the zone of albitization. c) Third type: post-magmatic deposits which are connected with ultra-basic alkali rocks. Carbonatites are mineralized; pyrochlore and hatchettolite as well as carbonates and fluocarbonates of the rare earths are found. The following radioactive minerals are found: thorianite, monazite, and zircon. The anomaly decreases in the direction from the center to the periphery of the intrusion. Granite pegmatites show only local accumulations of uranium and thorium minerals. Pneumatolytic-hydrothermal deposits could be observed by the enrichment of uranium and thorium. Some

Card 2/3

Prospecting For Rare and Disperse Elements
With Radioactive Methods

S/172/60/013/003-4/001/002
B002/B067

general remarks are made on the radiometric determination. Finally, it is said that the following deposits have already been determined by means of radiometric methods: diamond, bauxite, phosphorite, titanium, copper-bearing sandstones, black shale, and others. There are 7 references:
6 Soviet.

SUBMITTED: April 15, 1960

Card 3/3

LISTENGARTEN, B.M.; TER-OGANESOVA, A.G.

Determining reservoir rock permeability on the basis of the
study of water influx of wells. Azerb.neft.khoz. 35 no.5:6-8
My '56. (MLRA 9:10)

(Permeability) (Oil field flooding)

TER-OGANESOVA, V.S.

Conference of young scientists. Neft. khoz. 43 no.8:66 Ag
'65. (MIRA 18:12)

TER-OGANESYAN, G.N.

Perinephric novocaine block in complex treatment of tuberculous
mesoadenitis. Izv. AN Arm. SSR, Biol.nauki 12 no.10:89-94 0 '59.
(MIRA 13:3)

1. Khirurgicheskoye otdeleniye Dilizhanskoy gorodskoy bol'nitsy.
(MESENTERY--TUBERCULOSIS) (NOVOCAIN)

TER-OGANESYAN, G. N., Cand Med Sci -- (diss) "Perinephric novocainol Blockade in the System of the Complex Treatment of Tubercular mezadenites." Yerevan, 1960, 42 pages. (Yerevan State Med Inst), 200 copies. (KL, 29-60, 127)

TER-OGANESYAN, I. M.		PROCESSES AND PROPERTIES INDEX	
SA		B 64 H	
<p>621.116.915</p> <p>3319. Method of testing neutralizing devices on electrical equipment without disconnecting them from the network. I. M. Ter-Oganessian. <i>From: Energ.</i>, No. 3, 7-11 (March, 1951) in Russian.</p> <p>One of the phases of tested equipment is connected to its "earthed" frame through an additional resistance, current through which is measured together with voltage drop in the circuit, using an ammeter, voltmeter and a bridge. Claims of high accuracy, based on theoretical considerations, have been upheld by field tests.</p> <p>I. LUKASZEWICZ</p>			
<p>ASB-SLA METALLURGICAL LITERATURE CLASSIFICATION</p> <p>REGIONAL SYMBOLS</p> <p>SEARCHED INDEXED</p> <p>REVIEWED</p> <p>RECEIVED</p> <p>DATE</p> <p>BY</p> <p>INITIALS</p> <p>DATE</p> <p>BY</p> <p>INITIALS</p>			

TER-OGANESYAN, I. M.

"Methods of Testing the Zeroing Circuit of the Electric Equipment of Industrial Enterprises." Min Higher Education, Leningrad Electrotechnical Inst imeni V. I. Ul'yanov (Lenin), Leningrad, 1955. (Dissertation for the Degree of Candidate of Technical Sciences)

SO: M-972, 20 Feb 56